



## **PROPER PERFORMANCE OF AUTONOMIC FUNCTION TESTING**

### **Introduction**

Autonomic testing is a type of electrodiagnostic (EDX) medicine and clinical neurophysiology testing which can be used to assess sympathetic and parasympathetic neural pathways. While nerve conduction studies and needle electromyography provide information regarding large diameter myelinated fibers, the focus of autonomic testing is small diameter myelinated and unmyelinated fibers. By studying the function of these fiber pathways, autonomic testing is designed to evaluate a spectrum of clinical scenarios including small fiber autonomic involvement in suspected peripheral neuropathy, conditions associated with orthostatic intolerance, putative autonomic dysfunction in extrapyramidal and other central nervous system syndromes, paroxysmal conditions which may result from episodic abnormalities of the autonomic nervous system, and related disorders. Autonomic testing in EDX medicine may be employed as part of multidisciplinary evaluations of autonomic function which may include collaborative evaluations with colleagues in a wide range of medical specialties such as ophthalmology, cardiology, endocrinology, gastroenterology, internal medicine, nuclear medicine, urology, and others.

### **Qualifications and Training**

It is the AANEM's position that, in order to perform and interpret diagnostic studies of the autonomic nervous system appropriately and ensure quality patient care, the individual performing the studies must be a physician with special training in the diagnosis and treatment of disorders of the autonomic nervous system and in the application of particular neurophysiological techniques to study these disorders. The AANEM believes that the physician should have completed formal residency training in neurology, internal medicine, physical medicine & rehabilitation, or pediatrics and also be prepared to demonstrate appropriate clinical expertise in autonomic disorders. Such training should include detailed medical education on:

- the anatomical organization and functions of the central autonomic control centers;
- the parasympathetic, sympathetic, and enteric nervous systems;
- the basis of autonomic neurotransmission;
- the pharmacology of drugs affecting the autonomic nervous system;
- clinical and laboratory diagnosis of central and peripheral disorders of autonomic function;
- the indications, techniques, and limitations of noninvasive recordings of arterial pressure, heart rate, and sudomotor output including theoretical and practical knowledge of the equipment used to obtain those recordings; and
- the performance and interpretation of tests of orthostatic tolerance, adrenergic, cardiovagal, and sudomotor function.

Evidence of such expertise can be demonstrated in several ways, such as (1) subspecialty board certification in autonomic disorders by the United Council for Neurologic Subspecialties, (2) completion of 12 months of fellowship training in autonomic disorders, or (3) obtaining at least 50 hours of Accreditation Council for Continuing Medical Education approved category 1 CME specifically related to autonomic disorders plus substantial clinical practice in the diagnosis and treatment of patients with autonomic disorders.

### **Appropriate Performance**

The necessary steps for an appropriate EDX evaluation using autonomic testing is as follows:

Development of a differential diagnosis by a trained physician, based upon an appropriate history and physical examination performed by this physician. Once a physician has determined the preliminary differential diagnosis, a technician may perform the tests selected by the physician. It is the position of the AANEM that supervision by a physician, as defined by Medicare, is the minimum standard recommended for all autonomic testing.

Techniques used to study autonomic function include specific tests to evaluate cardiovagal, vasomotor adrenergic and sudomotor function. Cardiovagal testing can provide an early marker of autonomic parasympathetic dysfunction. Tests of sympathetic adrenergic function are the primary method for evaluating patients with syncope, orthostatic hypotension, postural orthostatic tachycardia syndrome, and postural dizziness. Sudomotor tests assess the integrity of pathways that control sweat function. For additional information on the appropriate performance, review the recent statement by the American Academy of Neurology on Autonomic Testing.<sup>1</sup>

It is the AANEM's position that a standard battery of autonomic tests (including tests of cardiovagal, sympathetic adrenergic and sudomotor function) are often used to investigate the integrity of the autonomic nervous system in patients with suspected autonomic disorders. However, the exact battery of tests may vary from patient to patient depending on the clinical scenario. Tests that assess the pathways in question are recommended.

The AANEM recommends the following series of tests as reliable and reproducible.

- For evaluation of sudomotor function: quantitative sudomotor axon reflex testing, thermoregulatory sweat testing, induced silastic skin imprints, and the sympathetic skin response.
- For evaluation of cardiovagal function: the heart rate response to deep breathing, Valsalva ratio, and postural change.
- For evaluation of sympathetic adrenergic function: the continuous beat-to-beat heart rate and blood pressure response to a Valsalva maneuver, tilt table test, or active standing.

The AANEM does not recommend pharmacologic testing to induce changes in blood pressure. Additional changes in monitored systolic or diastolic blood pressure and heart rate must be appropriately reviewed as obtained (on-site), prior to dismissing the patient. This step is necessary to determine if test repetition is required for any reasons including technical concerns and also to determine what other diagnostic tests are required.

Before results can be interpreted as normal or abnormal, it is important that the physician consider other factors that can cause an abnormal result including the beat-to-beat blood pressure setting, room temperature, ambient humidity, tobacco, caffeine or medication use, or any other improper setting including the

participation of the patient. The physician interpreting the studies should understand the factors that could cause abnormalities and interpret them in the context of the test when determining a diagnosis.

### **Summary**

The AANEM strongly recommends that autonomic testing procedures be performed by physicians with comprehensive knowledge of neurological and autonomic disorders to ensure precise interpretation and diagnosis at the completion of the testing.

Individuals without sufficient medical education in autonomic disorders are unqualified to interpret the data generated or to coordinate the findings with other clinical information to reach a clinical diagnosis. Subspecialists that perform organ specific tests in their own subspecialty should be well versed with autonomic pathways and basic principles of EDX medicine/electrophysiology.

It is also AANEM's position that the same physician should directly supervise and interpret the data on-site and in real time collected in various autonomic procedures including those performed by a technician.

**Approved by the American Association of Neuromuscular & Electrodiagnostic Medicine: August 2016.  
Revised and re-approved May 2017.**

### **REFERENCES**

1. American Academy of Neurology Autonomic Testing Model Coverage Policy.  
<https://www.aan.com/practice/billing-and-coding/coverage-policies/>