

# Report Sample

Helping physicians improve the quality of their reports is a key goal for the AANEM. The report template below is based on the AANEM's educational paper Reporting the Results of Nerve Conduction Studies and Needle EMG. A report template helps the EDX physician adhere to and document required procedures —by checking them off a list. This process will help the EDX physician complete a thorough analysis of the patient's

history, physical, and EDX data that will improve diagnostic accuracy and result in better quality patient care. The template also will help laboratories applying for the laboratory accreditation program meet the criteria used to evaluate EDX reports. The template was developed listing the key elements for an EDX standard report excluding F-wave, H-reflex, and repetitive stimulation studies. Physicians are strongly urged to utilize this template to improve their reports.

Patient Name: Patient Address:	Patient ID: 01234 Gender: Female Date of Birth: 01/01/65 Height/Weight: 5'7" 200lbs BMI: 31.3	Patient Demographics																																																																																																																																																																																																				
Lab Name, address, phone number: Referring Physician: Dr. X	Examination Date: 01/10/11	Reason for Referral																																																																																																																																																																																																				
<p><b>History and Physical Examination</b>  A 46-year-old right-handed female with a past medical history of right carpal tunnel syndrome status post release surgery and remote breast cancer status post right-sided mastectomy was referred for an EDX examination for worsening nocturnal and positionally provoked numbness and tingling in her left hand over the last 3 years. Symptoms are relieved with shaking out her hand. Pertinent medical records were reviewed, including the documented physical examination, and a focused history was obtained to confirm the clinical problem and reason for referral. The nature and purpose of the procedure were explained, the planned examination sites were confirmed with the patient, and a pre-procedure pause was performed to verify all safety elements before testing.</p> <p>These EDX studies were performed to evaluate for a suspected left carpal tunnel syndrome (CTS).</p> <p><b>NCS Examination</b>  For sensory nerve conduction studies, the amplitude is measured peak-to-peak, the latency reported is the peak latency, and the conduction velocity, if measured, is determined from onset latencies and is calculated over the forearm.  For motor nerve conduction studies, the amplitude is measured baseline-to-peak, the latency reported is the onset latency, and the conduction velocity is calculated over the forearm.</p> <p>Limb temperature: 32.8° C</p>																																																																																																																																																																																																						
<p><b>NERVE CONDUCTION STUDIES</b> [*Abnormal results; <sup>§</sup>AANEM Normative Data Task Force reference values (where applicable)]</p> <table border="1"> <thead> <tr> <th>Type of Study</th> <th>Side/ Nerve</th> <th>Stimulation Site</th> <th>Recording Site</th> <th>Distance<sup>§</sup></th> <th>AMP</th> <th>Reference Value<sup>§</sup></th> <th>Latency</th> <th>Reference Value<sup>§</sup></th> <th>Conduction Velocity</th> <th>Reference Value<sup>§</sup></th> </tr> </thead> <tbody> <tr> <td>Sensory</td> <td>Left Median</td> <td>Wrist</td> <td>Index</td> <td>14cm</td> <td>14<math>\mu</math>V</td> <td>(&gt;13 <math>\mu</math>V)</td> <td>3.9ms</td> <td>(&lt;4.0 ms)</td> <td></td> <td></td> </tr> <tr> <td>Mixed</td> <td>Left Median</td> <td>Palm</td> <td>Wrist</td> <td>7cm</td> <td>55<math>\mu</math>V</td> <td>(&gt;8 <math>\mu</math>V)</td> <td>2.5ms*</td> <td>(&lt;2.3 ms)</td> <td></td> <td></td> </tr> <tr> <td>Motor</td> <td>Left Median</td> <td>Wrist</td> <td>APB</td> <td>8cm</td> <td>13.5mV</td> <td>(&gt;4.2 mV)</td> <td>4.1ms</td> <td>(&lt;4.4ms)</td> <td></td> <td></td> </tr> <tr> <td></td> <td>Left Median</td> <td>Elbow/Wrist</td> <td>APB</td> <td>24cm</td> <td>12.9mV</td> <td>(&gt;4.2 mV)</td> <td>8.3ms</td> <td></td> <td>57 m/s</td> <td>(&gt;51m/s)</td> </tr> <tr> <td>Sensory</td> <td>Left Ulnar</td> <td>Wrist</td> <td>Digit V</td> <td>14cm</td> <td>12<math>\mu</math>V</td> <td>(&gt;8 <math>\mu</math>V)</td> <td>3.4ms</td> <td>(&lt;4.0 ms)</td> <td></td> <td></td> </tr> <tr> <td>Mixed</td> <td>Left Ulnar</td> <td>Palm</td> <td>Wrist</td> <td>8cm</td> <td>25<math>\mu</math>V</td> <td>(&gt;12 <math>\mu</math>V)</td> <td>2.0ms</td> <td>(&lt;2.3 ms)</td> <td></td> <td></td> </tr> <tr> <td>Motor</td> <td>Left Ulnar</td> <td>Wrist</td> <td>ADM</td> <td>8cm</td> <td>8.2mV</td> <td>(&gt;7.9 mV)</td> <td>3.1ms</td> <td>(&lt;3.7 ms)</td> <td></td> <td></td> </tr> <tr> <td></td> <td>Left Ulnar</td> <td>Elbow/wrist</td> <td>ADM</td> <td>21cm</td> <td>8.0mV</td> <td>(&gt;7.9 mV)</td> <td>6.9ms</td> <td></td> <td>55m/s</td> <td>(&gt;52 m/s)</td> </tr> </tbody> </table> <p><b>EMG Examination</b>  The study was performed with a concentric needle electrode.</p> <p><b>Needle EMG Results</b></p> <table border="1"> <thead> <tr> <th rowspan="2">Side</th> <th rowspan="2">Muscle</th> <th rowspan="2">Insertional Activity</th> <th colspan="3">Spontaneous Activity</th> <th colspan="4">Voluntary Activity</th> </tr> <tr> <th>Positive Sharp Waves</th> <th>Fibrillation</th> <th>Fasciculation</th> <th>Amplitude</th> <th>Duration</th> <th>%Polyphasics</th> <th>Recruitment</th> </tr> </thead> <tbody> <tr> <td>Left</td> <td>Deltoid</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Biceps</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Triceps</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Pronator Teres</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Extensor Indicis</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Flexor Pollicis Longus</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>First Dorsal Interosseous</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> <tr> <td>Left</td> <td>Abductor Pollicis Brevis</td> <td>Normal</td> <td>0</td> <td>0</td> <td>0</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> <td>Normal</td> </tr> </tbody> </table> <p><b>Findings</b></p> <ul style="list-style-type: none"> <li>The left median mixed nerve conduction study was ABNORMAL: The peak latency was prolonged with orthodromic stimulation of the palm.</li> <li>The left median sensory nerve conduction study revealed normal peak latency and amplitude.</li> <li>The left median motor nerve conduction study revealed normal onset latency, amplitude and conduction velocity.</li> <li>The left ulnar sensory and mixed nerve conduction studies revealed normal peak latencies and amplitudes.</li> <li>The left ulnar motor nerve conduction study revealed normal onset latency, amplitude and conduction velocity with elbow flexed at 90 degrees.</li> <li>Needle examination with a concentric needle electrode of selected muscles of the left upper extremity was normal.</li> </ul> <p><b>Diagnostic Interpretation</b></p> <ol style="list-style-type: none"> <li>The results of the study were ABNORMAL.</li> <li>The findings were compatible with a very mild left median mononeuropathy at or distal to the wrist, consistent with a clinical diagnosis of carpal tunnel syndrome.</li> <li>There was no electrodiagnostic evidence of a more proximal left median nerve pathology, left ulnar mononeuropathy, or left cervical motor radiculopathy in the areas tested (left C5-C8/T1-innervated muscles).</li> </ol> <p><b>Notes:</b>  Right upper limb comparison studies were not performed due to lymphedema following right mastectomy.  In comparison with a prior normal EDX study of the left upper limb done on 02/15/08, today's study demonstrates interval development of the left median neuropathy, as mentioned in #2 above.</p> <p><b>Physician Signature:</b>  EDX Physician, MD, 1/10/11, 2:00 PM</p>			Type of Study	Side/ Nerve	Stimulation Site	Recording Site	Distance <sup>§</sup>	AMP	Reference Value <sup>§</sup>	Latency	Reference Value <sup>§</sup>	Conduction Velocity	Reference Value <sup>§</sup>	Sensory	Left Median	Wrist	Index	14cm	14 $\mu$ V	(>13 $\mu$ V)	3.9ms	(<4.0 ms)			Mixed	Left Median	Palm	Wrist	7cm	55 $\mu$ V	(>8 $\mu$ V)	2.5ms*	(<2.3 ms)			Motor	Left Median	Wrist	APB	8cm	13.5mV	(>4.2 mV)	4.1ms	(<4.4ms)				Left Median	Elbow/Wrist	APB	24cm	12.9mV	(>4.2 mV)	8.3ms		57 m/s	(>51m/s)	Sensory	Left Ulnar	Wrist	Digit V	14cm	12 $\mu$ V	(>8 $\mu$ V)	3.4ms	(<4.0 ms)			Mixed	Left Ulnar	Palm	Wrist	8cm	25 $\mu$ V	(>12 $\mu$ V)	2.0ms	(<2.3 ms)			Motor	Left Ulnar	Wrist	ADM	8cm	8.2mV	(>7.9 mV)	3.1ms	(<3.7 ms)				Left Ulnar	Elbow/wrist	ADM	21cm	8.0mV	(>7.9 mV)	6.9ms		55m/s	(>52 m/s)	Side	Muscle	Insertional Activity	Spontaneous Activity			Voluntary Activity				Positive Sharp Waves	Fibrillation	Fasciculation	Amplitude	Duration	%Polyphasics	Recruitment	Left	Deltoid	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Biceps	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Triceps	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Pronator Teres	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Extensor Indicis	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Flexor Pollicis Longus	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	First Dorsal Interosseous	Normal	0	0	0	Normal	Normal	Normal	Normal	Left	Abductor Pollicis Brevis	Normal	0	0	0	Normal	Normal	Normal	Normal
Type of Study	Side/ Nerve	Stimulation Site	Recording Site	Distance <sup>§</sup>	AMP	Reference Value <sup>§</sup>	Latency	Reference Value <sup>§</sup>	Conduction Velocity	Reference Value <sup>§</sup>																																																																																																																																																																																												
Sensory	Left Median	Wrist	Index	14cm	14 $\mu$ V	(>13 $\mu$ V)	3.9ms	(<4.0 ms)																																																																																																																																																																																														
Mixed	Left Median	Palm	Wrist	7cm	55 $\mu$ V	(>8 $\mu$ V)	2.5ms*	(<2.3 ms)																																																																																																																																																																																														
Motor	Left Median	Wrist	APB	8cm	13.5mV	(>4.2 mV)	4.1ms	(<4.4ms)																																																																																																																																																																																														
	Left Median	Elbow/Wrist	APB	24cm	12.9mV	(>4.2 mV)	8.3ms		57 m/s	(>51m/s)																																																																																																																																																																																												
Sensory	Left Ulnar	Wrist	Digit V	14cm	12 $\mu$ V	(>8 $\mu$ V)	3.4ms	(<4.0 ms)																																																																																																																																																																																														
Mixed	Left Ulnar	Palm	Wrist	8cm	25 $\mu$ V	(>12 $\mu$ V)	2.0ms	(<2.3 ms)																																																																																																																																																																																														
Motor	Left Ulnar	Wrist	ADM	8cm	8.2mV	(>7.9 mV)	3.1ms	(<3.7 ms)																																																																																																																																																																																														
	Left Ulnar	Elbow/wrist	ADM	21cm	8.0mV	(>7.9 mV)	6.9ms		55m/s	(>52 m/s)																																																																																																																																																																																												
Side	Muscle	Insertional Activity	Spontaneous Activity			Voluntary Activity																																																																																																																																																																																																
			Positive Sharp Waves	Fibrillation	Fasciculation	Amplitude	Duration	%Polyphasics	Recruitment																																																																																																																																																																																													
Left	Deltoid	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Biceps	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Triceps	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Pronator Teres	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Extensor Indicis	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Flexor Pollicis Longus	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	First Dorsal Interosseous	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
Left	Abductor Pollicis Brevis	Normal	0	0	0	Normal	Normal	Normal	Normal																																																																																																																																																																																													
<p><b>Patient Demographics</b></p> <p><b>Reason for Referral</b></p> <p><b>Description of History and Physical Exam</b></p> <p><b>Limb Temperature</b>  &gt;32°C UE, &gt;30°C LE</p> <p><b>Tabular NCS Data:</b></p> <ul style="list-style-type: none"> <li>Side &amp; Nerve</li> <li>Stimulation &amp; Recording Site</li> <li>SNAP/CMAP Amplitude</li> <li>Distal Peak or Onset Latency</li> <li>Conduction Velocity</li> </ul> <p><b>Reference Values</b></p> <p><b>Tabular EMG Data:</b></p> <ul style="list-style-type: none"> <li>Side</li> <li>Muscle Tested</li> <li>Activity Data</li> <li>Voluntary</li> <li>Insertional</li> <li>Spontaneous</li> </ul> <p><b>Description of Findings</b></p> <p><b>Probable Diagnosis &amp; Location of Pathology</b></p> <p><b>Study Limitations &amp; Previous Study Information</b></p> <p><b>Signature with Time Stamp</b></p>																																																																																																																																																																																																						